

Brugada Syndrome Panel

Panel Gene List: *ABCC9, CACNA1C, CACNA2D1, CACNB2, GPD1L, HCN4, KCND3, KCNE3, KCNH2, KCNJ8, PKP2, SCN10A, SCN1B, SCN2B, SCN3B, SCN5A, TRPM4*

Additional genes from our cardiology test menu may be added to this panel by selecting test code 481C.

Clinical Features:

Brugada syndrome (BrS) is a genetic heart disorder due to abnormal ion channel function characterized by ST segment elevation on ECG (leads V1-3) in the absence of structural heart disease.¹⁻³ It is associated with increased risk for syncope, ventricular tachyarrhythmia and sudden cardiac death. In individuals with an apparently normal heart, Brugada syndrome accounts for up to 20% of unexpected sudden deaths and is suspected to account for 4-12% of all unexpected sudden deaths.⁴ Brugada syndrome occurs worldwide and is estimated to affect 5 per 10,000 individuals of all ethnicities, with some regional differences.³

The diagnosis of BrS is based on clinical history, ECG findings, and family history. Typically, the disorder manifests in patients between ages 20 to 40, but symptoms have been reported from infancy through late life. Most individuals with BrS are asymptomatic. The most common clinical symptoms are syncope and cardiac arrest that occur at rest, during sleep, or with high fever. In some patients, symptoms of BrS will develop after taking certain medications such as sodium channel blockers. Sudden cardiac death may occur without preceding symptoms and without an identifiable cause at autopsy. Additionally, many symptoms of BrS are similar to those of other heart conditions, such as arrhythmogenic right ventricular cardiomyopathy (ARVC), atypical right bundle branch block, left ventricular hypertrophy, early repolarization, acute myocardial infarction, and acute pericarditis.

Inheritance Pattern/Genetics: Autosomal Dominant

Test Methods:

Using genomic DNA from the submitted specimen, the complete coding regions and splice site junctions of the genes on this panel are enriched using a proprietary targeted capture system developed by GeneDx for next-generation sequencing with CNV calling (NGS-CNV) (Only exons 1-44 for CACNA1C and excluding exon 6 of the PKP2 gene). The enriched targets are simultaneously sequenced with paired-end reads on an Illumina platform. Bi-directional sequence reads are assembled and aligned to reference sequences based on NCBI RefSeq transcripts and human genome build GRCh37/UCSC hg19. After gene specific filtering, data are analyzed to identify sequence variants and most deletions and duplications involving coding exons. Alternative sequencing or copy number detection methods are used to analyze regions with inadequate sequence or copy number data. Reported clinically significant variants are confirmed by an appropriate orthogonal method. Reportable variants include pathogenic variants, likely pathogenic variants and variants of uncertain significance. Likely benign and benign variants, if present, are

not routinely reported but are available upon request. Sequencing and deletion/duplication analysis of the remaining genes on the Arrhythmia Panel is available as a separate test if the BrS Panel is negative.

Test Sensitivity:

The clinical sensitivity of sequencing and deletion/duplication analysis of the genes included in the BrS Panel depends in part on the patient's clinical phenotype and family history. In general, the sensitivity is highest for individuals with clearly defined BrS and a family history of disease. The technical sensitivity of sequencing is estimated to be > 99% at detecting single nucleotide events. It will not reliably detect deletions greater than 20 base pairs, insertions or rearrangements greater than 10 base pairs, or low-level mosaicism. The copy number assessment methods used with this test cannot reliably detect copy number variants of less than 500 base pairs or mosaicism and cannot identify balanced chromosome aberrations. Assessment of exon-level copy number events is dependent on the inherent sequence properties of the targeted regions, including shared homology and exon size. Gene specific exclusions for exon-level deletion/duplication testing for this panel are: SCN1B only whole gene deletions or duplications may be detected.

Gene	Protein	Inheritance	Disease Association(s)
<i>ABCC9</i>	ATP-BINDING CASSETTE, SUBFAMILY C, MEMBER 9	AD	DCM, BrS, ERS, Cantu syndrome and related disorders
<i>CACNA1C</i>	CALCIUM CHANNEL, VOLTAGE-DEPENDENT, L TYPE, ALPHA-1C SUBUNIT	AD	BrS (with short QTc), Timothy syndrome, LQTS
<i>CACNA2D1</i>	CALCIUM CHANNEL, VOLTAGE-DEPENDENT, ALPHA-2/DELTA SUBUNIT 1	AD	BrS, Epilepsy
<i>CACNB2</i>	CALCIUM CHANNEL, VOLTAGE-DEPENDENT, BETA-2 SUBUNIT	AD	BrS (with short QTc)
<i>GPD1L</i>	GLYCEROL-3-PHOSPHATE DEHYDROGENASE 1-LIKE	AD	BrS
<i>HCN4</i>	HYPERPOLARIZATION-ACTIVATED CYCLIC NUCLEOTIDE-GATED POTASSIUM CHANNEL 4	AD	BrS, SSS, AF, AV block, Bradycardia, Tachycardia, LVNC
<i>KCND3</i>	POTASSIUM CHANNEL, VOLTAGE-GATED, SHAL-RELATED SUBFAMILY, MEMBER 3	AD	BrS, AF, SIDS, Spinocerebellar ataxia
<i>KCNH2 (HERG)</i>	POTASSIUM CHANNEL, VOLTAGE-GATED, SUBFAMILY H, MEMBER 2	AD	BrS, LQTS, SQTS
<i>KCNE3</i>	POTASSIUM CHANNEL, VOLTAGE-GATED, ISK-RELATED SUBFAMILY, MEMBER 3	AD	BrS
<i>KCNJ8</i>	POTASSIUM CHANNEL, INWARDLY RECTIFYING, SUBFAMILY J, MEMBER 8	AD	BrS, VF, SIDS, Cantu syndrome
<i>PKP2</i>	PLAKOPHILIN 2	AD	ARVC, BrS
<i>SCN1B</i>	SODIUM CHANNEL, VOLTAGE-GATED, TYPE I, BETA SUBUNIT	AD	BrS, CCD, Epilepsy
<i>SCN2B</i>	SODIUM CHANNEL, VOLTAGE-GATED, TYPE II, BETA SUBUNIT	AD	BrS, AF
<i>SCN3B</i>	SODIUM CHANNEL, VOLTAGE-GATED, TYPE III, BETA SUBUNIT	AD	BrS, AF, VF, SIDS
<i>SCN5A</i>	SODIUM CHANNEL, VOLTAGE-GATED, TYPE V, ALPHA SUBUNIT	AD, AR	BrS, ARVC/ARVC-like disease, DCM, HB, LQTS, SIDS, SSS
<i>SCN10A</i>	SODIUM CHANNEL, VOLTAGE-GATED, TYPE X, ALPHA SUBUNIT	AD	BrS, LQTS, AF, painful small-fiber peripheral neuropathy
<i>TRPM4</i>	TRANSIENT RECEPTOR POTENTIAL CATION CHANNEL, SUBFAMILY M, MEMBER 4	AD	HB, BrS

Abbreviations: AD – Autosomal dominant; AF – Atrial fibrillation; AR – Autosomal recessive; ARVC- Arrhythmogenic right ventricular cardiomyopathy; AV – Atrioventricular; BrS – Brugada syndrome; CCD – Cardiac conduction defect; DCM – Dilated cardiomyopathy; ERS – Early repolarization syndrome; HB – Heart block; LQTS – Long QT syndrome; SIDS – Sudden infant death syndrome; SSS – Sick sinus syndrome; SIDS – Sudden infant death syndrome; VF – Ventricular fibrillation

References:

1. Brugada R, Campuzano O, Brugada P, et al. Brugada Syndrome. 2005 Mar 31 [Updated 2014 Apr 10]. In: Pagon RA, Adam MP, Ardinger HH, et al., editors. GeneReviews [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2014. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK1517/>
2. Hedley et al. (2009) Human Mutation 30 (9):1256-66 (PMID: 19606473)
3. Fowler et al. (2009) Current Opinion In Cardiology 24 (1):74-81 (PMID: 19102039)
4. Antzelevitch et al. (2002) Circ. Res. 91 (12):1114-8 (PMID: 12480811)