

AR Gene Analysis in Androgen Insensitivity Syndrome (AIS) / Sequencing and Deletion/Duplication

Disorder also known as: Testicular Feminization syndrome (TFM); Reifenstein syndrome

Clinical Features:

Androgen insensitivity syndrome may be complete or partial. Patients with AIS may come to attention in utero or at birth because of inconsistency between prenatal karyotype (male) and ultrasound findings of a female fetus, or at birth because of ambiguous genitalia. Alternatively, patients may present during the pubertal years because of a presumed inguinal hernia (abdominal or inguinal testes), absence of pubic/auxiliary hair, or lack of onset of menses. The mature phenotype is often characterized by well-developed breasts and luxuriant scalp hair. In the partial form, patients may exhibit hypospadias, micropenis, or fusion of the labial folds and undergo virilization at puberty. 46,XX individuals who are heterozygous carriers of an AR variant typically do not exhibit any clinical differences in sexual differentiation, although they may have patchy changes in hair distribution and irregular menses due to skewed X chromosome inactivation.

Of note, Kennedy disease is an independent disorder caused by an expansion of a CAG repeat in the AR gene and is not diagnosed with this test.

Genetics:

Androgen insensitivity syndrome (AIS) has an X-linked recessive pattern of inheritance.

Test Sensitivity:

Approximately 95-97% of individuals with AIS are expected to have a variant in the AR gene identifiable by sequencing, while 3-5% will have a deletion or duplication.⁹ The detection rate for individuals with milder phenotypes (i.e. partial androgen insensitivity and mild androgen insensitivity) is not well established but is likely less than 50%.^{1,2} Additionally, 5-6% of males with hypospadias have been found to harbor an identifiable variant in the AR gene.^{3,4,5}

Variant Spectrum:

Many distinct variants scattered across the AR gene have been identified in both complete and partial androgen insensitivity syndrome. The vast majority of the variants are missense substitutions, although nonsense and splice-site variants, whole and partial gene deletions, and two partial gene duplications also have been reported.^{8,1,6,7}

Test Methods:

Using genomic DNA from the submitted specimen, the coding regions and splice junctions of the AR gene are PCR amplified and capillary sequencing is performed. Bi-directional sequence is assembled, aligned to reference gene sequences based on NCBI RefSeq transcript and human genome build GRCh37/UCSC hg19, and is analyzed for sequence variants. Concurrent deletion/duplication testing is performed for most, if not all, of the coding exons using exon-level oligo array CGH (ExonArrayDx). Probe sequences and locations were based on human genome build GRCh37/UCSC hg19. In 46,XY individuals, deletions of one or more exons would be detectable by sequencing. The concurrent ExonArrayDx evaluates for smaller deletions as well as duplications in 46,XY individuals and deletions and duplications in 46,XX individuals. Reported clinically significant variants are confirmed by an appropriate method. Sequence and copy number alterations are reported according to the Human Genome Variation Society (HGVS) or International System for Human Cytogenetic Nomenclature (ISCN) guidelines, respectively. Reportable variants include pathogenic variants, likely pathogenic variants and variants of uncertain significance. Likely benign and benign variants are not routinely reported but are available upon request.

References:

1. Ahmed et al., (2000) JCEM. 85:658-665.
2. Batch et al., (1992) Hum Molec Genet 1:497-503.
3. Hiort et al., (1994) Eur J Pediatr 153:317-321.
4. Albers et al., (1997) J Pediatr 131:386-392.
5. Wang et al., (2004) Eur J Hum Genet 12:706-712.
6. Avila et al., (2002) JCEM 87:182-188.
7. Hannema et al., (2004) JCEM 89:5815-5822.
8. Gottlieb et al (2004) Hum Mutat 23:527-533.
9. Gottlieb B, Trifiro MA. Androgen Insensitivity Syndrome. 1999 Mar 24 [Updated 2017 May 11]. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2018.